

## HUMAN FGF GENE AND GENE EXPRESSION PRODUCTS

## REFERENCE TO RELATED APPLICATIONS

This application is a continuation of provisional U.S. Patent Application Serial No. 60/077,411, filed March 9, 1998, and U.S. Patent Application Serial No. 60/083,553, filed April 29, 1998.

INSAI

## TECHNICAL FIELD

The present invention relates to novel nucleic acid sequences encoding a member of the fibroblast growth factor (FGF) family, and to polypeptides encoded by the nucleic acid sequences.

## BACKGROUND OF THE INVENTION

This invention provides a gene of the fibroblast growth factor (FGF) family and protein encoded by the gene. The invention is directed to mRNA expressed by human cells, polynucleotides having coding regions corresponding to the mRNA, protein products of the polynucleotides and polypeptide products of mRNA, and biological function of the polypeptides and proteins.

At least 17 members of the FGF family have been identified, and are discussed in, for example, Ohuchi et al., *Development* 124:2235-2244 (1994); Ghosh et al., *Cell Growth and Differentiation* 7:1425-1434 (1996); Gemel et al., *Genomics* 35:253-257 (1996); Crossley et al., *Development* 121:439-451 (1995); and Crossley et al., *Cell* 84:127-136 (1996).

## SUMMARY OF THE INVENTION

The present invention provides a composition comprising an isolated polynucleotide selected from the group consisting of:

- (a) a polynucleotide comprising at least eight contiguous nucleotides from nucleotide 550 to 1170 of SEQ ID NO:4;
- (b) a polynucleotide that encodes a variant of the polypeptide encoded by (a); and

- (c) a polynucleotide encoding a protein expressed by a polynucleotide having the sequence of SEQ ID NO:4.

The invention further provides for the use of the isolated polynucleotides or fragments thereof as diagnostic probes or as primers.

The present invention also provides a composition comprising a polypeptide, wherein said polypeptide is selected from the group consisting of:

- (a) a polypeptide comprising at least 6 contiguous amino acids encoded by SEQ ID NO:4 in the region of nucleotide 550 to 1170;
- (b) a polypeptide encoded by a polynucleotide comprising SEQ ID NO:4; and
- (c) a variant of the protein (a) or (b).

In certain preferred embodiments of the invention, the polynucleotide is operably linked to an expression control sequence. The invention further provides a host cell, including bacterial, yeast, insect and mammalian cells, transformed with the polynucleotide sequence. The invention also provides full-length cDNA and full-length polynucleotides corresponding to SEQ ID NO:4.

Protein and polypeptide compositions of the invention may further comprise a pharmaceutically acceptable carrier. Compositions comprising an antibody that specifically reacts with such protein or polypeptide are also provided by the present invention.

The invention provides methods of using the FGF of the invention for the isolation, regeneration, proliferation, and differentiation of mammalian multipotent neural stem cells, progenitor cells and progeny.

In a preferred embodiment, primary central (CNS) and peripheral nervous system (PNS) cells when treated with FGF 98 of the invention proliferate, have at least a limited self regeneration capacity, and can undergo lineage restriction in response to the local environment.

In a further embodiment, cells produced by treatment with FGF 98 are used to screen drugs such as small molecules, and growth factors, which may affect development, differentiation, survival and or function of CNS and PNS derived neurons and glia.

The invention also provides for the production of large amounts of otherwise minor cell populations of cells to be used for generation of cDNA libraries for the isolation of rare molecules expressed in the precursors cells or progeny; cells produced by treatment may directly express growth factors or other molecules, and conditioned media is screened in assays for novel activities.

The invention further provides for the isolation, self-renewal and survival of mammalian neural stem cells and the differentiation of their progeny.

#### BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1. Western analysis of FGF 98 produced by SF-9 cells infected with different clones of baculovirus expressing FGF 98 (Clone 1 to 6). The highest producers were Clones 2 and 4.

Figure 2. (A) Elution profile of the CM of SF-9 cells expressing FGF 98 on a Heparin Sepharose (HS) affinity column. Figure 2(B) shows the OD profile of the 0.5M, 1M and 2M NaCl (● - ●) fractions together with the bioactivity profile of the various fractions (○) when tested on granulosa cells.

Figure 3. (A) shows SDS-PAGE analysis of the 2M NaCl fractions followed by Coomassie blue staining, indicating the presence of a single band with a MW of 28 kDa in the peak fractions of the 2M NaCl eluate (frac. 13, 14). Figure 3(B) shows Western analysis using an anti-peptide FGF 98 antibody of the HS fractions shown in Figure 2A. A single band with a MW of 28 kD is seen in fractions 12 to 16 corresponding to the 2M NaCl fractions.

Figure 4. The active HS fractions (Fractions 2-16) were loaded onto a C4 Vydac column (25x0.46 cm, 5  $\mu$ m particle size, 300 Å pore size) equilibrated in 0.1% (v/v) TFA, 20% acetonitrile. Protein was eluted with a 115 min. gradient of 20-100% acetonitrile in 0.1% TFA at a flow rate of 0.6 ml/min. at room temperature. Fractions of 1.2 ml were collected. Aliquots of each fraction were diluted 1 to 10 with 0.2% gelatin in PBS and bioassayed on granulosa cells. A major peak of protein closely associated with a small peak could be observed. Both protein peaks were bioactive.

Figure 5. (A) SDS-PAGE of the RP-HPLC fractions shown in Figure 4. When stained by Coomassie blue, both Fractions 22 and 23 gave a single band with a MW of

28 kDa. Figure 5(B). When analyzed by Western using an FGF 98 anti-peptide antibody both Fractions 22 and 23 were recognized as a single band with a MW of 28 kDa.

Figure 6. Heparin Sepharose chromatography of a 3 liter batch of medium conditioned by SF-9 cells expressing FGF 98. The CM was adjusted to 0.5M NaCl and pH to 7.3. It was applied on a column of Heparin Sepharose equilibrated in 0.5M NaCl 10 mM Tris pH 7.2 on sequentially eluted with 1M and 2M NaCl (A, B).

Figure 7. (A) SDS-PAGE of the 1M and 2M NaCl fractions of Figure 6 followed by Coomassie blue staining. Figure 7(B). Western analysis of the 1M and 2M NaCl fractions of Figure 6 using a specific FGF 98 antipeptide antibody.

Figure 8. Mono S ion-exchange chromatography of the HS-purified and active fractions. The bioactive fractions eluted from HS were affinity pooled and diluted 10-fold with 10 mM Tris pH 7.2. Using a 50 ml Super loop, the sample was then applied on a Mono S HR 5/5 column equilibrated in 10 mM Tris pH 7.2-0.2 M NaCl at room temperature. The column was eluted with a multilinear gradient of NaCl (0.2M to 2M NaCl). Absorbency was monitored at 280 nm. Flow rate was 1 ml/min and 1 ml fractions were collected. Aliquots of each fraction were diluted 1 to 100 in 0.2% gelatin in PBS, and 10  $\mu$ l aliquots were bioassayed on granulosa cells in 12 well cluster plates as described.

Figure 9. Comparison of the biological activity for granulosa cells of FGF 98 (●) and basic FGF(0).

Figure 10 is a graph showing the effect of FGF 98 on neural stem cell proliferation. Dotted bars represent 6 days *in vitro* (DIV) and solid bars represent 21 days *in vitro* (DIV). The treatment consisted of no additions (1), bFGF (2), or FGF 98 of the invention (3).

Figure 11. Panel A is a photograph of bFGF-treated rat primary hippocampal cell cultures, showing some nestin expression. Panel B shows a separate culture of cells treated with FGF 98 of the invention. More extensive nestin expression is seen.

Figure 12. Panel A shows FGF 98 mRNA expression in adult brain stem and cerebellum. Panel B shows a high magnification of FGF 98 mRNA expression in motor neurons of the facial nucleus.

Fig 2A-B  
Fig 3A-B  
Fig 5A-B  
Fig 7A-B  
Fig 11  
Purified  
not  
labeled

## DETAILED DESCRIPTION OF THE INVENTION

Because of their potent activities for promoting growth, proliferation, survival and differentiation of a wide variety of cells and tissue types, FGFs continue to be pursued as therapeutic agents for a number of different indications, including wound healing, such as musculo-skeletal conditions, for example, bone fractures, ligament and tissue repair, tendonitis, bursitis, etc.; skin conditions, for example, burns, cuts, lacerations, bed sores, slow healing ulcers, etc.; tissue protection and repair during myocardial infarction and ischemia, in the treatment of neurological conditions, for example, neuro-degenerative disease and stroke, in the treatment of eye disease, including macular degeneration, and the like.

The fibroblast growth factor (FGF) proteins identified to date belong to a family of signaling molecules that regulate growth and differentiation of a variety of cell types. The significance of FGF proteins to human physiology and pathology relates in part to their key roles in embryogenesis, in blood vessel development and growth, and in bone growth. *In vitro* experiments have demonstrated a role for FGF in regulating cell growth and division of endothelial cells, vascular smooth muscle cells, fibroblasts, and cardiac and skeletal myocytes. Other members of the FGF family and their biological roles are described in Crossley et al., *Development* 121:439-451 (1995); Ohuchi et al., *Development* 124:2235-2244 (1997); Gemel et al., *Genomics* 35:253-257 (1996); and Ghosh et al., *Cell Growth and Differentiation* 7:1425-1434 (1996).

FGF proteins are also significant to human health and disease because of a role in cancer cell growth. For example, FGF-8 was identified as an androgen-induced growth factor in breast and prostate cancer cells. (Tanaka et al., *FEBS Lett.* 363:226-230 (1995) and *P.N.A.S.* 89:8928-8932 (1992)).

A member of the FGF family is described here, wherein the FGF protein is expressed at high levels in adult mouse brain, and plays a protective role in neuronal tissue. A polynucleotide encoding the FGF of the invention has the sequence as shown in SEQ ID NO:1. The polynucleotide was identified as encoding a member of the FGF family by the conserved regions throughout the amino acid sequence and by the regions of homology shared by the polynucleotide and genes encoding known FGF proteins.

The FGF of the invention shares greatest homology with FGF-8, which plays a variety of roles, including outgrowth and patterning in mouse embryo development (Crossley et al., *Development* 121:439-451 (1995), and midbrain development (Crossley et al., *Nature* 380:66-68 (1996). Eight FGF-8 isoforms have been identified in mice, generated by alternative splicing, and may be functionally redundant. FGF-8b is 100% identical in mice and humans. Blunt et al., *J. Biol. Chem.* 272:3733-3738 (1997). FGF-8 and FGF-10 function in a coordinate manner during chick limb development. Ohuchi et al., *Development* 124:2235-2244 (1997).

As the above brief discussion illustrates, the knowledge about the roles played by various FGF proteins continues to grow, but is by far incomplete.

The present invention adds to this knowledge by disclosing that the FGF of SEQ ID NO:5 plays a role in adult mouse brain and has a protective role in neuronal tissue. The FGF 98 increased the number of neuronal cells at six and twenty days *in vitro*. Furthermore, unlike bFGF-treated cells, few FGF 98-treated cells die under the conditions used herein. FGF 98 can be used for *ex vivo* expansion of precursor cells in the nervous system. Transplant or engraftment of the cells can be used as a therapy for diseases in which specific populations of neurons are lost due to degeneration such as, for example, Parkinson's disease and Alzheimer's disease.

The invention therefore is based upon the identification, isolation and sequencing of a new fibroblast growth factor (FGF 98). This growth factor has been found to promote cell survival and, in particular, the survival of neurons.

Reference to FGF 98 herein is intended to be construed to include growth factors of any origin which are substantially homologous to and which are biologically equivalent to the FGF 98 characterized and described herein. Such substantially homologous growth factors may be native to any tissue or species and, similarly, biological activity can be characterized in any of a number of biological assay systems.

The terms "biologically equivalent" are intended to mean that the compositions of the present invention are capable of demonstrating some or all of the same growth properties in a similar fashion, not necessarily to the same degree as the FGF 98 isolated as described herein or recombinantly produced human FGF 98 of the invention.

By "substantially homologous" it is meant that the degree of homology of human FGF 98 to FGF 98 from any species is greater than that between FGF 98 and any previously reported member of the FGF family.

Sequence identity or percent identity is intended to mean the percentage of same residues between two sequences, referenced to human FGF when determining percent identity with non-human FGF 98, referenced to FGF 98 when determining percent identity with non-FGF 98 growth factors, when the two sequences are aligned using the Clustal method (Higgins et al, *Cabios* 8:189-191, 1992) of multiple sequence alignment in the Lasergene biocomputing software (DNASTAR, INC, Madison, WI). In this method, multiple alignments are carried out in a progressive manner, in which larger and larger alignment groups are assembled using similarity scores calculated from a series of pairwise alignments. Optimal sequence alignments are obtained by finding the maximum alignment score, which is the average of all scores between the separate residues in the alignment, determined from a residue weight table representing the probability of a given amino acid change occurring in two related proteins over a given evolutionary interval. Penalties for opening and lengthening gaps in the alignment contribute to the score. The default parameters used with this program are as follows: gap penalty for multiple alignment=10; gap length penalty for multiple alignment=10; k-tuple value in pairwise alignment=1; gap penalty in pairwise alignment=3; window value in pairwise alignment=5; diagonals saved in pairwise alignment=5. The residue weight table used for the alignment program is PAM250 (Dayhoff et al., in *Atlas of Protein Sequence and Structure*, Dayhoff, Ed., NDRF, Washington, Vol. 5, suppl. 3, p. 345, 1978).

Percent conservation is calculated from the above alignment by adding the percentage of identical residues to the percentage of positions at which the two residues represent a conservative substitution (defined as having a log odds value of greater than or equal to 0.3 in the PAM250 residue weight table). Conservation is referenced to human FGF 98 when determining percent conservation with non-human FGF 98, referenced to FGF 98 when determining percent conservation with non-FGF 98 growth factors. Conservative amino acid changes satisfying this requirement are: R-K; E-D, Y-F, L-M; V-I, Q-H.

FGF 98 can also include hybrid and modified forms of FGF 98 including fusion proteins and FGF 98 fragments and hybrid and modified forms in which certain amino acids

have been deleted or replaced and modifications such as where one or more amino acids have been changed to a modified amino acid or unusual amino acid and modifications such as glycosylations so long as the hybrid or modified form retains the biological activity of FGF 98. By retaining the biological activity, it is meant that neuronal survival is promoted, although not necessarily at the same level of potency as that of the FGF 98 isolated as described herein or that of the recombinantly produced human FGF 98.

Also included within the meaning of substantially homologous is any FGF 98 which may be isolated by virtue of cross-reactivity with antibodies to the FGF 98 described herein or whose encoding nucleotide sequences including genomic DNA, mRNA or cDNA may be isolated through hybridization with the complementary sequence of genomic or subgenomic nucleotide sequences or cDNA of the FGF 98 herein or fragments thereof. It will also be appreciated by one skilled in the art that degenerate DNA sequences can encode human FGF 98 and these are also intended to be included within the present invention as are allelic variants of FGF 98.

Growth factors are thought to act at specific receptors. According to the invention, FGF 98 and as yet unknown members of this family of growth factors act through specific receptors having distinct distributions as has been shown for other growth factor families.

A preferred FGF 98 of the present invention has been identified and isolated in purified form as described. Also preferred is FGF 98 prepared by recombinant DNA technology. By "pure form" or "purified form" or "substantially purified form" it is meant that an FGF 98 composition is substantially free of other proteins which are not FGF 98.

Recombinant human FGF 98 may be made by expressing the DNA sequences encoding FGF 98 in a suitable transformed host cell. Using methods well known in the art, the DNA encoding FGF 98 may be linked to an expression vector, transformed into a host cell and conditions established that are suitable for expression of FGF 98 by the transformed cell.

Any suitable expression vector may be employed to produce recombinant human FGF 98 such as expression vectors for use in insect cells. Baculovirus expression systems can also be employed. A preferable method is expression in insect cells, such as Tr5 or Sf9 cells, using baculovirus vector, as described in Example 1.



The present invention includes nucleic acid sequences including sequences that encode human FGF 98. Also included within the scope of this invention are sequences that are substantially the same as the nucleic acid sequences encoding FGF 98. Such substantially the same sequences may, for example, be substituted with codons more readily expressed in a given host cell such as *E. coli* according to well known and standard procedures. Such modified nucleic acid sequences are included within the scope of this invention.

Specific nucleic acid sequences can be modified by those skilled in the art and, thus, all nucleic acid sequences that code for the amino acid sequences of FGF 98 can likewise be so modified. The present invention thus also includes nucleic acid sequence which will hybridize with all such nucleic acid sequences--or complements of the nucleic acid sequences where appropriate--and encode a polypeptide having the cell survival promoting activities disclosed herein. The present invention also includes nucleic acid sequences that encode for polypeptides that have neuronal survival promoting activity and that are recognized by antibodies that bind to FGF 98. A preferred method for raising antibody is described in Example 9.

The present invention also encompasses vectors comprising expression regulatory elements operably linked to any of the nucleic acid sequences included within the scope of the invention. This invention also includes host cells--of any variety--that have been transformed with vectors comprising expression regulatory elements operably linked to any of the nucleic acid sequences included within the scope of the present invention.

Methods are also provided herein for producing FGF 98. Preparation can be by isolation from conditioned medium from a variety of cell types so long as the cell type produces FGF 98. A second and preferred method involves utilization of recombinant methods by isolating or obtaining a nucleic acid sequence encoding FGF 98, cloning the sequence along with appropriate regulatory sequences into suitable vectors and cell types, and expressing the sequence to produce FGF 98.

Although FGF 98 has been described on the basis of its ability to promote the survival of particular cells, this factor will act on other neuronal cell types as well. For example, FGF 98 is shown herein to enhance nestin expression. Thus, it is likely that FGF 98 can act on a wide variety of central and peripheral neuronal cell types.

It is also likely that FGF 98 will act on non-neuronal cells to promote their survival, growth or function. This expectation is based upon the activity of known growth factors. Members of the FGF family act on many cell types of different function and embryologic origin.

The inventors herein have identified several non-neuronal tissues in which FGF 98 is expressed including lung, heart, developing bone, and chondrocytes. This suggests a role for FGF 98 in, for example, maintenance of musculoskeletal function and treatment of related disease; correction of congenital defects; and treatment cardiomyopathies.

The present invention also includes therapeutic or pharmaceutical compositions comprising FGF 98 in an effective amount for treating patients with cellular degeneration and a method comprising administering a therapeutically effective amount of FGF 98. These compositions and methods are useful for treating a number of degenerative diseases. Where the cellular degeneration involves neuronal degeneration, the diseases include, but are not limited to peripheral neuropathy, amyotrophic lateral sclerosis, Alzheimer's disease, Parkinson's disease, Huntington's disease, ischemic stroke, acute brain injury, acute spinal chord injury, nervous system tumors, multiple sclerosis, peripheral nerve trauma or injury, exposure to neurotoxins, metabolic diseases such as diabetes or renal dysfunctions and damage caused by infectious agents. Where the cellular degeneration involves bone marrow cell degeneration, the diseases include, but are not limited to disorders of insufficient blood cells such as, for example, leukopenias including eosinopenia and/or basopenia, lymphopenia, monocytopenia, neutropenia, anemias, thrombocytopenia as well as an insufficiency of stem cells for any of the above. The above cells and tissues can also be treated for depressed function.

The compositions and methods herein can also be useful to prevent degeneration and/or promote survival in other non-neuronal tissues as well. One skilled in the art can readily determine using a variety of assays known in the art for identifying whether FGF 98 would be useful in promoting survival or functioning in a particular cell type, such as cardiac cells.

In certain circumstances, it may be desirable to modulate or decrease the amount of FGF 98 expressed. Thus, in another aspect of the present invention, FGF 98 anti-

sense oligonucleotides can be made and a method utilized for diminishing the level of expression of FGF 98 by a cell comprising administering one or more FGF 98 anti-sense oligonucleotides. By FGF 98 anti-sense oligonucleotides reference is made to oligonucleotides that have a nucleotide sequence that interacts through base pairing with a specific complementary nucleic acid sequence involved in the expression of FGF 98 such that the expression of FGF 98 is reduced. Preferably, the specific nucleic acid sequence involved in the expression of FGF 98 is a genomic DNA molecule or mRNA molecule that encodes FGF 98. This genomic DNA molecule can comprise regulatory regions of the FGF 98 gene, or the coding sequence for mature FGF 98 protein. The term complementary to a nucleotide sequence in the context of FGF 98 antisense oligonucleotides and methods therefor means sufficiently complementary to such a sequence as to allow hybridization to that sequence in a cell, *i.e.*, under physiological conditions. The FGF 98 antisense oligonucleotides preferably comprise a sequence containing from about 8 to about 100 nucleotides and more preferably the FGF 98 antisense oligonucleotides comprise from about 15 to about 30 nucleotides. The FGF 98 antisense oligonucleotides can also contain a variety of modifications that confer resistance to nucleolytic degradation such as, for example, modified internucleoside linkages (Uhlmann and Peyman, *Chemical Reviews* 90:543-548 1990; Schneider and Banner, *Tetrahedron Lett.* 31:335, 1990 which are incorporated by reference), modified nucleic acid bases and/or sugars and the like.

The therapeutic or pharmaceutical compositions of the present invention can be administered by any suitable route known in the art including for example intravenous, subcutaneous, intramuscular, transdermal, intrathecal or intracerebral. Administration can be either rapid as by injection or over a period of time as by slow infusion or administration of slow release formulation. For treating tissues in the central nervous system, administration can be by injection or infusion into the cerebrospinal fluid (CSF). When it is intended that FGF 98 be administered to cells in the central nervous system, administration can be with one or more agents capable of promoting penetration of FGF 98 across the blood-brain barrier.

FGF 98 can also be linked or conjugated with agents that provide desirable pharmaceutical or pharmacodynamic properties. For example, FGF 98 can be coupled to any substance known in the art to promote penetration or transport across the blood-brain barrier

such as an antibody to the transferrin receptor, and administered by intravenous injection (See, for example, Friden et al., *Science* 259:373-377, 1993 which is incorporated by reference). Furthermore, FGF 98 can be stably linked to a polymer such as polyethylene glycol to obtain desirable properties of solubility, stability, half-life and other pharmaceutically advantageous properties. (See, for example, Davis et al., *Enzyme Eng.* 4:169-73, 1978; Burnham, *Am. J. Hosp. Pharm.* 51:210-218, 1994 which are incorporated by reference).

The compositions are usually employed in the form of pharmaceutical preparations. Such preparations are made in a manner well known in the pharmaceutical art. One preferred preparation utilizes a vehicle of physiological saline solution, but it is contemplated that other pharmaceutically acceptable carriers such as physiological concentrations of other non-toxic salts, five percent aqueous glucose solution, sterile water or the like may also be used. It may also be desirable that a suitable buffer be present in the composition. Such solutions can, if desired, be lyophilized and stored in a sterile ampoule ready for reconstitution by the addition of sterile water for ready injection. The primary solvent can be aqueous or alternatively non-aqueous. FGF 98 can also be incorporated into a solid or semi-solid biologically compatible matrix which can be implanted into tissues requiring treatment.

The carrier can also contain other pharmaceutically-acceptable excipients for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the carrier may contain still other pharmaceutically-acceptable excipients for modifying or maintaining release or absorption or penetration across the blood-brain barrier. Such excipients are those substances usually and customarily employed to formulate dosages for parenteral administration in either unit dosage or multi-dose form or for direct infusion into the cerebrospinal fluid by continuous or periodic infusion.

Dose administration can be repeated depending upon the pharmacokinetic parameters of the dosage formulation and the route of administration used.

It is also contemplated that certain formulations containing FGF 98 are to be administered orally. Such formulations are preferably encapsulated and formulated with suitable carriers in solid dosage forms. Some examples of suitable carriers, excipients, and

diluents include lactose, dextrose, sucrose, sorbitol, mannitol, starches, gum acacia, calcium phosphate, alginates, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, cellulose, gelatin, syrup, methyl cellulose, methyl- and propylhydroxybenzoates, talc, magnesium, stearate, water, mineral oil, and the like. The formulations can additionally include lubricating agents, wetting agents, emulsifying and suspending agents, preserving agents, sweetening agents or flavoring agents. The compositions may be formulated so as to provide rapid, sustained, or delayed release of the active ingredients after administration to the patient by employing procedures well known in the art. The formulations can also contain substances that diminish proteolytic degradation and promote absorption such as, for example, surface active agents.

The specific dose is calculated according to the approximate body weight or body surface area of the patient or the volume of body space to be occupied. The dose will also be calculated dependent upon the particular route of administration selected. Further refinement of the calculations necessary to determine the appropriate dosage for treatment is routinely made by those of ordinary skill in the art. Such calculations can be made without undue experimentation by one skilled in the art in light of the activity disclosed herein in assay preparations of target cells. Exact dosages are determined in conjunction with standard dose-response studies. It will be understood that the amount of the composition actually administered will be determined by a practitioner, in the light of the relevant circumstances including the condition or conditions to be treated, the choice of composition to be administered, the age, weight, and response of the individual patient, the severity of the patient's symptoms, and the chosen route of administration.

In one embodiment of this invention, FGF 98 may be therapeutically administered by implanting into patients vectors or cells capable of producing a biologically-active form of FGF 98 or a precursor of FGF 98, *i.e.*, a molecule that can be readily converted to a biological-active form of FGF 98 by the body. In one approach cells that secrete FGF 98 may be encapsulated into semipermeable membranes for implantation into a patient. The cells can be cells that normally express FGF 98 or a precursor thereof or the cells can be transformed to express FGF 98 or a precursor thereof. It is preferred that the cell be of human origin and that the FGF 98 be human FGF 98 when the patient is human. However, the

formulations and methods herein can be used for veterinary as well as human applications and the term "patient" as used herein is intended to include human and veterinary patients.

Cells can be grown *ex vivo* for use in transplantation or engraftment into patients (Muench et al., *Leuk. & Lymph.* 16:1-11, 1994 which is incorporated by reference). In another embodiment of the present invention, FGF 98 is used to promote the *ex vivo* expansion of a cells for transplantation or engraftment. Current methods have used bioreactor culture systems containing factors such as erythropoietin, colony stimulating factors, stem cell factor, and interleukins to expand hematopoietic progenitor cells for erythrocytes, monocytes, neutrophils, and lymphocytes (Verfaillie, *Stem Cells* 12:466-476, 1994 which is incorporated by reference). These stem cells can be isolated from the marrow of human donors, from human peripheral blood, or from umbilical cord blood cells. The expanded blood cells are used to treat patients who lack these cells as a result of specific disease conditions or as a result of high dose chemotherapy for treatment of malignancy (George, *Stem Cells* 12(Suppl 1):249-255, 1994 which is incorporated by reference). In the case of cell transplant after chemotherapy, autologous transplants can be performed by removing bone marrow cells before chemotherapy, expanding the cells *ex vivo* using methods that also function to purge malignant cells, and transplanting the expanded cells back into the patient following chemotherapy (for review, see Rummel and Van Zant, *J. Hematotherapy* 3:213-218, 1994 which is incorporated by reference). Since FGF 98 is expressed in the developing animal in blood, bone marrow and liver, tissues where proliferation and differentiation of progenitor cells occur, it is believed that FGF 98 can function to regulate the proliferation of hematopoietic stem cells and the differentiation of mature hematopoietic cells. Thus, the addition of FGF 98 to culture systems used for *ex vivo* expansion of cells could stimulate the rate at which certain populations of cells multiply or differentiate, and improve the effectiveness of these expansion systems in generating cells needed for transplant.

It is also believed that FGF 98 can be used for the *ex vivo* expansion of precursor cells in the nervous system. Transplant or engraftment of cells is currently being explored as a therapy for diseases in which certain populations of neurons are lost due to degeneration such as, for example, in Parkinson's disease (Bjorklund, *Curr. Opin. Neurobiol.* 2:683-689, 1992 which is incorporated by reference). Neuronal precursor cells can be

obtained from animal or human donors or from human fetal tissue and then expanded in culture using FGF 98 or other growth factors. These cells can then be engrafted into patients where they would function to replace some of the cells lost due to degeneration. Because neurotrophins have been shown to be capable of stimulating the survival and proliferation of neuronal precursors cells such as, for example, NT-3 stimulation of sympathetic neuroblast cells (Birren et al., *Develop.* 119:597-610, 1993 which is incorporated by reference), FGF 98 could also function in similar ways during the development of the nervous system and could be useful in the ex vivo expansion of neuronal cells.

In a number of circumstances it would be desirable to determine the levels of FGF 98 in a patient. The identification of FGF 98 along with the present report showing that FGF 98 is expressed by a number of tissues provides the basis for the conclusion that the presence of FGF 98 serves a normal physiologic function related to cell growth and survival. Indeed, other neurotrophic factors are known to play a role in the function of neuronal and non-neuronal tissues. (For review see Scully and Otten, *Cell Bol. Int.* 19:459-469, 1995; Otten and Gadiant, *Int. J. Devl. Neurosciences* 13:147-151, 1995 which are incorporated by reference). Endogenously produced FGF 98 may also play a role in certain disease conditions, particularly where there is cellular degeneration such as in neurodegenerative conditions or diseases. Other neurotrophic factors are known to change during disease conditions. For example, in multiple sclerosis, levels of NGF protein in the cerebrospinal fluid are increased during acute phases of the disease (Bracci-Laudiero et al., *Neuroscience Lett.* 147:9-12, 1992 which is incorporated by reference) and in systemic lupus erythematosus there is a correlation between inflammatory episodes and NGF levels in sera (Bracci-Laudiero et al. *NeuroReport* 4:563-565, 1993 which is incorporated by reference).

Given that FGF 98 is expressed in developing bone and in heart tissue, it is likely that the level of FGF 98 may be altered in a variety of conditions and that quantification of FGF 98 levels would provide clinically useful information. Furthermore, in the treatment of degenerative conditions, compositions containing FGF 98 can be administered and it would likely be desirable to achieve certain target levels of FGF 98 in sera, in cerebrospinal fluid or in any desired tissue compartment. It would, therefore, be advantageous to be able to monitor

the levels of FGF 98 in a patient. Accordingly, the present invention also provides methods for detecting the presence of FGF 98 in a sample from a patient.

The term "detection" as used herein in the context of detecting the presence of FGF 98 in a patient is intended to include the determining of the amount of FGF 98 or the ability to express an amount of FGF 98 in a patient, the distinguishing of FGF 98 from other growth factors, the estimation of prognosis in terms of probable outcome of a degenerative disease and prospect for recovery, the monitoring of the FGF 98 levels over a period of time as a measure of status of the condition, and the monitoring of FGF 98 levels for determining a preferred therapeutic regimen for the patient.

To detect the presence of FGF 98 in a patient, a sample is obtained from the patient. The sample can be a tissue biopsy sample or a sample of blood, plasma, serum, CSF or the like. FGF 98 is expressed in a variety of tissues as discussed in Example 8. Samples for detecting FGF 98 can be taken from any of these tissues. When assessing peripheral levels of FGF 98, it is preferred that the sample be a sample of blood, plasma or serum. When assessing the levels of FGF 98 in the central nervous system a preferred sample is a sample obtained from cerebrospinal fluid.

In some instances it is desirable to determine whether the FGF 98 gene is intact in the patient or in a tissue or cell line within the patient. By an intact FGF 98 gene it is meant that there are no alterations in the gene such as point mutations, deletions, insertions, chromosomal breakage, chromosomal rearrangements and the like wherein such alteration might alter production of FGF 98 or alter its biological activity, stability or the like to lead to disease processes or susceptibility to cellular degenerative conditions. Thus, in one embodiment of the present invention a method is provided for detecting and characterizing any alterations in the FGF 98 gene. The method comprises providing an oligonucleotide that contains the FGF 98 cDNA, genomic DNA or a fragment thereof or a derivative thereof. By a derivative of an oligonucleotide, it is meant that the derived oligonucleotide is substantially the same as the sequence from which it is derived in that the derived sequence has sufficient sequence complementarity to the sequence from which it is derived to hybridize to the FGF 98 gene. The derived nucleotide sequence is not necessarily physically derived from the



nucleotide sequence, but may be generated in any manner including for example, chemical synthesis or DNA replication or reverse transcription or transcription.

Typically, patient genomic DNA is isolated from a cell sample from the patient and digested with one or more restriction endonucleases such as, for example, TaqI and AluI. Using the Southern blot protocol, which is well known in the art, this assay determines whether a patient or a particular tissue in a patient has an intact FGF 98 gene or an FGF 98 gene abnormality.

Hybridization to an FGF 98 gene would involve denaturing the chromosomal DNA to obtain a single-stranded DNA; contacting the single-stranded DNA with a gene probe associated with the FGF 98 gene sequence; and identifying the hybridized DNA-probe to detect chromosomal DNA containing at least a portion of a human FGF 98 gene.

The term "probe" as used herein refers to a structure comprised of a polynucleotide that forms a hybrid structure with a target sequence, due to complementarity of probe sequence with a sequence in the target region. Oligomers suitable for use as probes may contain a minimum of about 8-12 contiguous nucleotides which are complementary to the targeted sequence and preferably a minimum of about 20.

The FGF 98 gene probes of the present invention can be DNA or RNA oligonucleotides and can be made by any method known in the art such as, for example, excision, transcription or chemical synthesis. Probes may be labeled with any detectable label known in the art such as, for example, radioactive or fluorescent labels or enzymatic marker. Labeling of the probe can be accomplished by any method known in the art such as by PCR, random priming, end labeling, nick translation or the like. One skilled in the art will also recognize that other methods not employing a labeled probe can be used to determine the hybridization. Examples of methods that can be used for detecting hybridization include Southern blotting, fluorescence in situ hybridization, and single-strand conformation polymorphism with PCR amplification.

Hybridization is typically carried out at 25° - 45° C, more preferably at 32° - 40° C and more preferably at 37° - 38° C. The time required for hybridization is from about 0.25 to about 96 hours, more preferably from about one to about 72 hours, and most preferably from about 4 to about 24 hours.

FGF 98 gene abnormalities can also be detected by using the PCR method and primers that flank or lie within the FGF 98 gene. The PCR method is well known in the art. Briefly, this method is performed using two oligonucleotide primers which are capable of hybridizing to the nucleic acid sequences flanking a target sequence that lies within an FGF 98 gene and amplifying the target sequence. The terms "oligonucleotide primer" as used herein refers to a short strand of DNA or RNA ranging in length from about 8 to about 30 bases. The upstream and downstream primers are typically from about 20 to about 30 base pairs in length and hybridize to the flanking regions for replication of the nucleotide sequence. The polymerization is catalyzed by a DNA-polymerase in the presence of deoxynucleotide triphosphates or nucleotide analogs to produce double-stranded DNA molecules. The double strands are then separated by any denaturing method including physical, chemical or enzymatic. Commonly, the method of physical denaturation is used involving heating the nucleic acid, typically to temperatures from about 80°C. to 105°C. for times ranging from about 1 to about 10 minutes. The process is repeated for the desired number of cycles.

The primers are selected to be substantially complementary to the strand of DNA being amplified. Therefore, the primers need not reflect the exact sequence of the template, but must be sufficiently complementary to selectively hybridize with the strand being amplified.

After PCR amplification, the DNA sequence comprising FGF 98 or pre-pro FGF 98 or a fragment thereof is then directly sequenced and analyzed by comparison of the sequence with the sequences disclosed herein to identify alterations which might change activity or expression levels or the like.

In another embodiment, a method for detecting FGF 98 is provided based upon an analysis of tissue expressing the FGF 98 gene. Certain tissues such as those identified below in Example 8 have been found to express the FGF 98 gene. The method comprises hybridizing a polynucleotide to mRNA from a sample of tissues that normally express the FGF 98 gene. The sample is obtained from a patient suspected of having an abnormality in the FGF 98 gene or in the FGF 98 gene of particular cells. The polynucleotide comprises SEQ ID NO:4 or a derivative thereof or a fragment thereof.

To detect the presence of mRNA encoding FGF 98 protein, a sample is obtained from a patient. The sample can be from blood or from a tissue biopsy sample. The sample may be treated to extract the nucleic acids contained therein. The resulting nucleic acid from the sample is subjected to gel electrophoresis or other size separation techniques.

The mRNA of the sample is contacted with a DNA sequence serving as a probe to form hybrid duplexes. The use of a labeled probes as discussed above allows detection of the resulting duplex.

When using the cDNA encoding FGF 98 protein or a derivative of the cDNA as a probe, high stringency conditions can be used in order to prevent false positives, that is the hybridization and apparent detection of FGF 98 nucleotide sequences when in fact an intact and functioning FGF 98 gene is not present. When using sequences derived from the FGF 98 cDNA, less stringent conditions could be used, however, this would be a less preferred approach because of the likelihood of false positives. The stringency of hybridization is determined by a number of factors during hybridization and during the washing procedure, including temperature, ionic strength, length of time and concentration of formamide. These factors are outlined in, for example, Sambrook et al. (Sambrook, et al., 1989, supra).

In order to increase the sensitivity of the detection in a sample of mRNA encoding the FGF 98 protein, the technique of reverse transcription/polymerization chain reaction (RT/PCR) can be used to amplify cDNA transcribed from mRNA encoding the FGF 98 protein. The method of RT/PCR is well known in the art (see example 9 and Figure 6 below).

The RT/PCR method can be performed as follows. Total cellular RNA is isolated by, for example, the standard guanidium isothiocyanate method and the total RNA is reverse transcribed. The reverse transcription method involves synthesis of DNA on a template of RNA using a reverse transcriptase enzyme and a 3' end primer. Typically, the primer contains an oligo(dT) sequence. The cDNA thus produced is then amplified using the PCR method and FGF 98 specific primers. (Belyavsky et al., *Nucl. Acid Res.* 17:2919-2932, 1989; Krug and Berger, *Methods in Enzymology*, 152:316-325, Academic Press, NY, 1987 which are incorporated by reference).

The polymerase chain reaction method is performed as described above using two oligonucleotide primers that are substantially complementary to the two flanking regions of the DNA segment to be amplified.

Following amplification, the PCR product is then electrophoresed and detected by ethidium bromide staining or by phosphoimaging.

The present invention further provides for methods to detect the presence of the FGF 98 protein in a sample obtained from a patient. Any method known in the art for detecting proteins can be used. Such methods include, but are not limited to immunodiffusion, immunoelectrophoresis, immunochemical methods, binder-ligand assays, immunohistochemical techniques, agglutination and complement assays. (for example, see *Basic and Clinical Immunology*, 217-262, Sites and Terr, eds., Appleton & Lange, Norwalk, CT, 1991 which is incorporated by reference). Preferred are binder-ligand immunoassay methods including reacting antibodies with an epitope or epitopes of the FGF 98 protein and competitively displacing a labeled FGF 98 protein or derivative thereof. A preferred antibody is prepared according to Example 9.

As used herein, a derivative of the FGF 98 protein is intended to include a polypeptide in which certain amino acids have been deleted or replaced or changed to modified or unusual amino acids wherein the FGF 98 derivative is biologically equivalent to FGF 98 and wherein the polypeptide derivative cross-reacts with antibodies raised against the FGF 98 protein. By cross-reaction it is meant that an antibody reacts with an antigen other than the one that induced its formation.

Numerous competitive and non-competitive protein binding immunoassays are well known in the art. Antibodies employed in such assays may be unlabeled, for example as used in agglutination tests, or labeled for use in a wide variety of assay methods. Labels that can be used include radionuclides, enzymes, fluorescers, chemilumescers, enzyme substrates or co-factors, enzyme inhibitors, particles, dyes and the like for use in radioimmunoassay (RIA), enzyme immunoassays, *e.g.*, enzyme-linked immunosorbent assay (ELISA), fluorescent immunoassays and the like.

Polyclonal or monoclonal antibodies to the FGF 98 protein or an epitope thereof can be made for use in immunoassays by any of a number of methods known in the art.

By epitope reference is made to an antigenic determinant of a polypeptide. An epitope could comprise 3 amino acids in a spacial conformation which is unique to the epitope. Generally an epitope consists of at least 5 such amino acids. Methods of determining the spatial conformation of amino acids are known in the art, and include, for example, x-ray crystallography and 2 dimensional nuclear magnetic resonance.

One approach for preparing antibodies to a protein is the selection and preparation of an amino acid sequence of all or part of the protein, chemically synthesizing the sequence and injecting it into an appropriate animal, usually a rabbit or a mouse (See Example 9).

Oligopeptides can be selected as candidates for the production of an antibody to the FGF 98 protein based upon the oligopeptides lying in hydrophilic regions, which are thus likely to be exposed in the mature protein. A preferred oligopeptide is KRYPKGQPELQKPFK (SEQ ID NO:6).

Antibodies to FGF 98 can also be raised against oligopeptides that include one or more of the conserved regions identified herein such that the antibody can cross-react with other family members. Such antibodies can be used to identify and isolate the other family members.

Methods for preparation of the FGF 98 protein or an epitope thereof include, but are not limited to chemical synthesis, recombinant DNA techniques or isolation from biological samples. Chemical synthesis of a peptide can be performed, for example, by the classical Merrifield method of solid phase peptide synthesis (Merrifield, *J. Am. Chem. Soc.* 85:2149, 1963 which is incorporated by reference) or the FMOC strategy on a Rapid Automated Multiple Peptide Synthesis system (E. I. du Pont de Nemours Company, Wilmington, DE) (Caprino and Han, *J Org Chem* 37:3404, 1972 which is incorporated by reference).

Polyclonal antibodies can be prepared by immunizing rabbits or other animals by injecting antigen followed by subsequent boosts at appropriate intervals. The animals are bled and sera assayed against purified FGF 98 protein usually by ELISA or by bioassay based upon the ability to block the action of FGF 98 on neurons or other cells. When using avian species, e.g., chicken, turkey and the like, the antibody can be isolated from the yolk of the

egg. Monoclonal antibodies can be prepared after the method of Milstein and Kohler by fusing splenocytes from immunized mice with continuously replicating tumor cells such as myeloma or lymphoma cells. (Milstein and Kohler, *Nature* 256:495-497, 1975; Galfre and Milstein, *Methods in Enzymology: Immunochemical Techniques* 73:1-46, Langone and Banatis eds., Academic Press, 1981 which are incorporated by reference). The hybridoma cells so formed are then cloned by limiting dilution methods and supernates assayed for antibody production by ELISA, RIA or bioassay.

The unique ability of antibodies to recognize and specifically bind to target proteins provides an approach for treating an over expression of the protein. Thus, another aspect of the present invention provides for a method for preventing or treating diseases involving over expression of the FGF 98 protein by treatment of a patient with specific antibodies to the FGF 98 protein.

Specific antibodies, either polyclonal or monoclonal, to the FGF 98 protein can be produced by any suitable method known in the art as discussed above. For example, murine or human monoclonal antibodies can be produced by hybridoma technology or, alternatively, the FGF 98 protein, or an immunologically active fragment thereof, or an anti-idiotypic antibody, or fragment thereof can be administered to an animal to elicit the production of antibodies capable of recognizing and binding to the FGF 98 protein. Such antibodies can be from any class of antibodies including, but not limited to IgG, IgA, IgM, IgD, and IgE or in the case of avian species, IgY and from any subclass of antibodies.

Polypeptides encoded by the instant polynucleotides and corresponding full-length genes can be used to screen peptide libraries, protein libraries, small molecule libraries, and phage display libraries, and other known methods, to identify analogs or antagonists.

Native FGF polypeptides may play a role in cancer. For example, FGF family members can induce marked morphological transformation of NIH 3T3 cells, and exhibit strong tumorigenicity in nude mice. Angiogenic activity has been exhibited by FGF family members. Thus, inhibitors of FGF can be used to treat cancer, such as prostate.

A library of peptides may be synthesized following the methods disclosed in U.S. Patent No. 5,010,175, and in PCT No. WO 91/17823. As described below in brief, a mixture of peptides is prepared, which is then screened to identify the peptides exhibiting the

desired signal transduction and receptor binding activity. According to the method of the '175 patent, a suitable peptide synthesis support (*e.g.*, a resin) is coupled to a mixture of appropriately protected, activated amino acids. The concentration of each amino acid in the reaction mixture is balanced or adjusted in inverse proportion to its coupling reaction rate so that the product is an equimolar mixture of amino acids coupled to the starting resin. The bound amino acids are then deprotected, and reacted with another balanced amino acid mixture to form an equimolar mixture of all possible dipeptides. This process is repeated until a mixture of peptides of the desired length (*e.g.*, hexamers) is formed. Note that one need not include all amino acids in each step: one may include only one or two amino acids in some steps (*e.g.*, where it is known that a particular amino acid is essential in a given position), thus reducing the complexity of the mixture. After the synthesis of the peptide library is completed, the mixture of peptides is screened for binding to the selected polypeptide. The peptides are then tested for their ability to inhibit or enhance activity. Peptides exhibiting the desired activity are then isolated and sequenced.

The method described in PCT No. WO 91/17823 is similar. However, instead of reacting the synthesis resin with a mixture of activated amino acids, the resin is divided into twenty equal portions (or into a number of portions corresponding to the number of different amino acids to be added in that step), and each amino acid is coupled individually to its portion of resin. The resin portions are then combined, mixed, and again divided into a number of equal portions for reaction with the second amino acid. In this manner, each reaction may be easily driven to completion. Additionally, one may maintain separate "subpools" by treating portions in parallel, rather than combining all resins at each step. This simplifies the process of determining which peptides are responsible for any observed receptor binding or signal transduction activity.

In such cases, the subpools containing, *e.g.*, 1-2,000 candidates each are exposed to one or more polypeptides of the invention. Each subpool that produces a positive result is then resynthesized as a group of smaller subpools (sub-subpools) containing, *e.g.*, 20-100 candidates, and reassayed. Positive sub-subpools may be resynthesized as individual compounds, and assayed finally to determine the peptides that exhibit a high binding constant. These peptides can be tested for their ability to inhibit or enhance the native activity. The

methods described in PCT No. WO 91/7823 and U.S. Patent No. 5,194,392 (herein incorporated by reference) enable the preparation of such pools and subpools by automated techniques in parallel, such that all synthesis and resynthesis may be performed in a matter of days.

Peptide agonists or antagonists are screened using any available method, such as signal transduction, antibody binding, receptor binding and mitogenic assays. The assay conditions ideally should resemble the conditions under which the native activity is exhibited *in vivo*, that is, under physiologic pH, temperature, and ionic strength. Suitable agonists or antagonists will exhibit strong inhibition or enhancement of the native activity at concentrations that do not cause toxic side effects in the subject. Agonists or antagonists that compete for binding to the native polypeptide may require concentrations equal to or greater than the native concentration, while inhibitors capable of binding irreversibly to the polypeptide may be added in concentrations on the order of the native concentration.

The end results of such screening and experimentation will be at least one novel polypeptide binding partner, such as a receptor, encoded by a gene or a cDNA of the invention, and at least one peptide agonist or antagonist of the novel binding partner. Such agonists and antagonists can be used to modulate, enhance, or inhibit receptor function in cells to which the receptor is native, or in cells that possess the receptor as a result of genetic engineering. Further, if the novel receptor shares biologically important characteristics with a known receptor, information about agonist/antagonist binding may help in developing improved agonists/antagonists of the known receptor.

The therapeutic FGF 98 polynucleotides and polypeptides of the present invention may be utilized in gene delivery vehicles. The gene delivery vehicle may be of viral or non-viral origin (*see generally*, Jolly, *Cancer Gene Therapy* 1:51-64 (1994); Kimura, *Human Gene Therapy* 5:845-852 (1994); Connelly, *Human Gene Therapy* 1:185-193 (1995); and Kaplitt, *Nature Genetics* 6:148-153 (1994)). Gene therapy vehicles for delivery of constructs including a coding sequence of a therapeutic of the invention can be administered either locally or systemically. These constructs can utilize viral or non-viral vector approaches. Expression of such coding sequences can be induced using endogenous



mammalian or heterologous promoters. Expression of the coding sequence can be either constitutive or regulated.

The present invention can employ recombinant retroviruses which are constructed to carry or express a selected nucleic acid molecule of interest. Retrovirus vectors that can be employed include those described in EP 0 415 731; WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 93/11230; WO 93/10218; Vile and Hart, *Cancer Res.* 53:3860-3864 (1993); Vile and Hart, *Cancer Res.* 53:962-967 (1993); Ram et al., *Cancer Res.* 53:83-88 (1993); Takamiya et al., *J. Neurosci. Res.* 33:493-503 (1992); Baba et al., *J. Neurosurg.* 79:729-735 (1993); U.S. Patent No. 4,777,127; GB Patent No. 2,200,651; and EP 0 345 242. Preferred recombinant retroviruses include those described in WO 91/02805.

Packaging cell lines suitable for use with the above-described retroviral vector constructs may be readily prepared (see PCT publications WO 95/30763 and WO 92/05266), and used to create producer cell lines (also termed vector cell lines) for the production of recombinant vector particles. Within particularly preferred embodiments of the invention, packaging cell lines are made from human (such as HT1080 cells) or mink parent cell lines, thereby allowing production of recombinant retroviruses that can survive inactivation in human serum.

The present invention also employs alphavirus-based vectors that can function as gene delivery vehicles. Such vectors can be constructed from a wide variety of alphaviruses, including, for example, Sindbis virus vectors, Semliki forest virus (ATCC VR-67; ATCC VR-1247), Ross River virus (ATCC VR-373; ATCC VR-1246) and Venezuelan equine encephalitis virus (ATCC VR-923; ATCC VR-1250; ATCC VR 1249; ATCC VR-532). Representative examples of such vector systems include those described in U.S. Patent Nos. 5,091,309; 5,217,879; and 5,185,440; and PCT Publication Nos. WO 92/10578; WO 94/21792; WO 95/27069; WO 95/27044; and WO 95/07994.

Gene delivery vehicles of the present invention can also employ parvovirus such as adeno-associated virus (AAV) vectors. Representative examples include the AAV vectors disclosed by Srivastava in WO 93/09239, Samulski et al., *J. Vir.* 63:3822-3828

(1989); Mendelson et al., *Virol.* 166:154-165 (1988); and Flotte et al., *P.N.A.S.* 90:10613-10617 (1993).

Representative examples of adenoviral vectors include those described by Berkner, *Biotechniques* 6:616-627 (Biotechniques); Rosenfeld et al., *Science* 252:431-434 (1991); WO 93/19191; Kolls et al., *P.N.A.S.* :215-219 (1994); Kass-Eisler et al., *P.N.A.S.* 90:11498-11502 (1993); Guzman et al., *Circulation* 88:2838-2848 (1993); Guzman et al., *Cir. Res.* 73:1202-1207 (1993); Zabner et al., *Cell* 75:207-216 (1993); Li et al., *Hum. Gene Ther.* 4:403-409 (1993); Cailaud et al., *Eur. J. Neurosci.* 5:1287-1291 (1993); Vincent et al., *Nat. Genet.* 5:130-134 (1993); Jaffe et al., *Nat. Genet.* 1:372-378 (1992); and Levrero et al., *Gene* 101:195-202 (1992). Exemplary adenoviral gene therapy vectors employable in this invention also include those described in WO 94/12649, WO 93/03769; WO 93/19191; WO 94/28938; WO 95/11984 and WO 95/00655. Administration of DNA linked to killed adenovirus as described in Curiel, *Hum. Gene Ther.* 3:147-154 (1992) may be employed.

Other gene delivery vehicles and methods may be employed, including polycationic condensed DNA linked or unlinked to killed adenovirus alone, for example Curiel, *Hum. Gene Ther.* 3:147-154 (1992); ligand-linked DNA, for example see Wu, *J. Biol. Chem.* 264:16985-16987 (1989); eukaryotic cell delivery vehicles cells, for example see U.S. Serial No. 08/240,030, filed May 9, 1994, and U.S. Serial No. 08/404,796; deposition of photopolymerized hydrogel materials; hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; ionizing radiation as described in U.S. Patent No. 5,206,152 and in WO 92/11033; nucleic charge neutralization or fusion with cell membranes. Additional approaches are described in Philip, *Mol. Cell Biol.* 14:2411-2418 (1994), and in Woffendin, *Proc. Natl. Acad. Sci.* 91:1581-1585 (1994).

Naked DNA may also be employed. Exemplary naked DNA introduction methods are described in WO 90/11092 and U.S. Patent No. 5,580,859. Uptake efficiency may be improved using biodegradable latex beads. DNA coated latex beads are efficiently transported into cells after endocytosis initiation by the beads. The method may be improved further by treatment of the beads to increase hydrophobicity and thereby facilitate disruption of the endosome and release of the DNA into the cytoplasm. Liposomes that can act as gene

delivery vehicles are described in U.S. Patent No. 5,422,120, PCT Patent Publication Nos. WO 95/13796, WO 94/23697, and WO 91/14445, and EP No. 0 524 968.

Further non-viral delivery suitable for use includes mechanical delivery systems such as the approach described in Woffendin et al., *Proc. Natl. Acad. Sci. USA* 91(24):11581-11585 (1994). Moreover, the coding sequence and the product of expression of such can be delivered through deposition of photopolymerized hydrogel materials. Other conventional methods for gene delivery that can be used for delivery of the coding sequence include, for example, use of hand-held gene transfer particle gun, as described in U.S. Patent No. 5,149,655; use of ionizing radiation for activating transferred gene, as described in U.S. Patent No. 5,206,152 and PCT Patent Publication No. WO 92/11033.

FGF has been implicated in diseases characterized by loss of function, inadequate function/number, abnormal function or death of cells, tissues or organs for which function or survival can be prolonged/rescued, and abnormalities reversed or prevented by therapy with FGF.

According to the invention, FGF 98 is likely to play a role in degenerative and other neurological diseases (disease characterized by loss of function or death of CNS cells, including neurons, glial cells, astrocytes, and others), in which therapy may prolong survival of cells or rescue/preserve function. Examples include Parkinson's disease, Alzheimers or other dementia, stroke, trauma, sequelae of infection, epilepsy, cranial nerve injury or dysfunction. FGF 98 may also be useful in treatment of eye disease characterized by similar responses of retinal cells, structural cells, nerve cells, etc., to administration of the molecule. Examples include Retinitis Pigmentosa, age-related macular degeneration, retinal detachment, ischemic retinal disease, etc.

FGF 98 may be useful in treatment of disorders of hearing, smell or taste due to loss/dysfunction of cells responsive to the molecule, including cranial or peripheral nerves, sensory cells, supporting cells, etc., and in treatment of peripheral motor or sensor nerve disease characterized by loss of function or death of neurons, nerves, other CNS or PNS cells or supporting cells: Amyotrophic Lateral Sclerosis, motor or Sensory Neuropathies (due to many causes; viral infections, metabolic disease-diabetes, autoimmune insult, idiopathic, degenerative, etc.) trauma.

Ischemic vascular disease may be amenable to FGF 98 treatment, wherein the disease is characterized by inadequate blood flow to an organ(s). Treatment may induce therapeutic angiogenesis or preserve function/survival of cells (myocardial ischemia/infarction, peripheral vascular disease, renal artery disease, stroke). Cardiomyopathies characterized by loss of function or death of cardiac myocytes or supporting cells in the heart (congestive heart failure, myocarditis) may also be treated using FGF 98, as can musculoskeletal disease characterized by loss of function, inadequate function or death of skeletal muscle cells, bone cells or supporting cells. Examples include skeletal myopathies, bone disease, and arthritis.

FGF 98 polynucleotides and polypeptides may aid in correction of congenital defects due to loss of FGF 98 molecule or its function (heart, lung, brain, limbs, kidney, etc.).

Treatment of wound healing is yet another use of FGF 98 polypeptides and polynucleotides, either due to trauma, disease, medical or surgical treatment, including regeneration of cell populations and tissues depleted by these processes. Examples include liver regeneration, operative wound healing, re-endothelialization of injured blood vessels, healing of traumatic wounds, healing of ulcers due to vascular, metabolic disease, etc., bone fractures, loss of cells due to inflammatory disease, etc.

FGF 98 may also be used in screens to identify drugs for treatment of cancers which involve over activity of the molecule, or new targets which would be useful in the identification of new drugs.

For all of the preceding embodiments, the clinician will determine, based on the specific condition, whether FGF 98 polypeptides or polynucleotides, antibodies to FGF 98, or small molecules such as peptide analogues or antagonists, will be the most suitable form of treatment. These forms are all within the scope of the invention.

Preferred embodiments of the invention are described in the following examples. Other embodiments within the scope of the claims herein will be apparent to one skilled in the art from consideration of the specification or practice of the invention as disclosed herein. It is intended that the specification, together with the examples, be considered exemplary only, with the scope and spirit of the invention being indicated by the claims which follow the examples.

## EXAMPLES

### EXAMPLE 1

#### EXPRESSION OF FGF 98 IN INSECT CELLS

The baculovirus vectors for expression of FGF 98 were generated by PCR of the coding region from a Bluescript vector containing the cDNA for FGF 98, prepared from a human heart cDNA library. Restriction sites for PstI and KpnI were added to facilitate directional subcloning into Bluebac 4.5 baculovirus vector (Invitrogen).

Supernatants from insect cells were generated for purification and bioassay by infecting SF9 cells at approximately  $1-2 \times 10^6$ /ml with recombinant baculovirus derived from these plasmids. A multiplicity of infection (moi) of 2-5 was used for the 48 hour infections. SF9 cells were removed from the supernatant by centrifugation and the supernatants further clarified by 0.2  $\mu$ M sterile filtration. In order to determine high producers, SF9 cells infected with different clones of baculovirus supernatants were analyzed by Western blot (Figure 1). A peptide antibody directed against the C terminal region of FGF 98 (AA177 to 195) was used. The yield in insect cells was 4 to 8 mg protein per liter of CM.

### EXAMPLE 2

#### ISOLATION OF FGF 98

Recombinant FGF 98 from the conditioned medium of SF-9 cells was purified by Heparin Sepharose (HS) affinity chromatography, followed by Mono S cation exchange chromatography. The recombinant FGF 98 bound to the HS was eluted using a step-wise salt gradient. Next, the recombinant FGF 98 eluted from the HS column was bound to the Mono S cation exchange column and was eluted using a nonlinear salt gradient.

Further details of the procedure utilized are described below.

#### A. Heparin Sepharose® Affinity Chromatography

The conditioned medium from Example 1 (30 ml) was loaded through a 5 ml bed of HS resin. The column was equilibrated in a buffer containing 150 mM NaCl and 10

mM Tris-HCl at pH 7.3. Once the conditioned medium was loaded, the column was washed extensively with the equilibration buffer until the absorbency at 280 nm returned to baseline. The protein was eluted from the HS column with an increasing step-wise NaCl gradient. The NaCl concentrations were 0.5M, 1M, and 2M NaCl, in 10 mM Tris-HCl at pH 7.3. The flow rate of the column during elution was approximately 30 ml/hr and 0.5-ml size fractions were collected.

The fractions were tested for FGF 98 bioactivity utilizing granulosa cells. The assay is described in Section 4 below. The fractions with the highest bioactivity were eluted with 2M NaCl (Figure 2).

When analyzed by SDS-PAGE followed by Coomassie blue staining the peak 2M NaCl fractions gave a single band with a MW of 28 kDa (Figure 3A). When analyzed by Western using an anti-FGF 98 peptide antibody the 2M NaCl fractions were immunoactive with a single band with a MW of 28 kDa (Figure 3B). No immunoreactive bands could be observed with the 1M NaCl fractions.

The homogeneity of the pooled 2M NaCl fractions was also analyzed by RP-HPLC. A single major protein peak eluting at 40% acetonitrile was observed. Closely associated with it was a minor protein peak that accounted for less than 5% of the protein mass (Figure 4). When the HPLC fractions were analyzed for bioactivity using granuloma cells, both protein peaks were active. When analyzed by SDS-PAGE followed by Coomassie blue staining, both fractions gave a single band with an apparent MW of 28k (Figure 5A). When this was followed by Western analysis both bands were recognized by a peptide antibody specific for FGF 98 (Figure 5B).

However, when the bioactivity of the pooled 2M NaCl fractions was analyzed on vascular endothelial cells, cytotoxicity was observed, indicating the presence of endotoxin-like substances. Those were eliminated using Mono S cation exchange chromatography as described below.

#### B. Large scale isolation of FGF 98.

A 3-liter batch of SF-9 cell-conditioned medium with pH adjusted to 7.3 and NaCl concentration raised to 0.5M was then allowed to run overnight at 4°C through a 30 ml

bed of HS resin. The column was equilibrated in a buffer containing 0.5M NaCl and 10 mM Tris- HCl pH 7.3. Following loading, the column was washed with the same buffer. The bound proteins were then eluted with 1M and 2M NaCl in 10 mM Tris pH 7.3. The flow rate of the column during the elution was 90 ml/hr and 3 ml size reactions were collected (Figures 6A, B).

When the collected fractions were tested for bioactivity using granulosa cells, all of the bioactivity was present in the 2M NaCl fractions. When the 1M and 2M NaCl fractions were analyzed by SDS-PAGE, the 1M NaCl fraction showed multiple bands with MW ranging from 200 kDa to 8 kDa. In contrast, the 2M NaCl fractions show a major single band with a MW of 28 kDa (Figure 7A). When analyzed by Western using an FGF 98 peptide antibody, the 1M NaCl contained a very weak immunoreactive band with a MW of 28 kDa. In contrast the 2M NaCl showed a very strong immunoreactive band at 28 kDa together with a weak band at 56 kDa (Figure 7B).

When the 2M NaCl fractions were tested on vascular endothelial cells, all fractions were toxic, indicating the presence of endotoxin-like material. Those were eliminated using Mono S cation exchange chromatography (Figure 8).

#### C. Mono S Cation Exchange Chromatography

The pooled 2M NaCl fractions eluted from the HS column were diluted ten-fold with 10 mM Tris-HCl at pH 7.3 to a final salt concentration of 0.2M NaCl. The material was then loaded with a Super loop onto a Mono S column linked to a FPLC system (Pharmacia, Piscataway, NJ). The Mono S cation exchange column was equilibrated with 10 mM Tris-HCl at pH 7.2. When the pooled fractions were loaded, the column was washed extensively at a flow rate of 1 ml/min. with the equilibration buffer until the absorbency returned to baseline. Then, protein was eluted from the column with a nonlinear NaCl gradient, 0.2M to 2M NaCl in 10 mM Tris-HCl at pH 7.3 at a flow rate of 1 ml/min. and 1 ml fractions were collected.

One major protein peak of activity was observed that eluted at about 0.8M NaCl (Figure 8). Fractions across the protein peak were assayed for bioactivity and subjected

to SDS-PAGE analysis. The results of the gel analysis (not illustrated) showed that protein from NaCl fractions exhibited an apparent molecular weight of 28 kD.

Western analysis of the Mono S 0.8M pooled fraction with peptide antibodies directed against AA176 to AA190 of FGF 98 revealed a single band with an apparent MW of 28 kD.

The homogeneity of the pooled Mono S fractions was also analyzed by RP-HPLC using a C4 column equilibrated in 0.1% TFA, 20% acetonitrile, using an acetonitrile gradient of 20 to 100%. A profile similar to that shown in Figure 4 was obtained (data not shown).

### EXAMPLE 3

#### FGF 98 BIOACTIVITY EFFECT ON GRANULOSA CELLS

Stock cultures of granulosa cells were grown and maintained in Dulbecco's modified Eagle medium supplemented with 10% bovine serum, 0.25  $\mu$ l/ml fungizone, and 10 ng/ml FGF. The cells were incubated at 37°C in a 10% CO<sub>2</sub> atmosphere with 99% humidity. For the bioactivity assay, the cells were seeded in 12-well plates at a density of  $5 \times 10^3$  cells per well in 1 ml of medium as described above for the stock cultures, and as described in Gospodarowicz et al. *Meth. Enzymol.* 147B:106-119 (D. Barnes et al., ed.) Academic Press, Orlando, FL (1987).

Ten microliter aliquots of the desired column fractions were diluted into 1 ml of 0.2% (w/v) gelatin in phosphate buffered saline (PBS). Ten microliters of this dilution were added to granulosa cells seeded in 12-well cluster plates containing 22 mm wells, at  $5 \times 10^3$  cells per well, and a 10  $\mu$ l aliquot of either the diluted column fractions or medium containing 10 ng bFGF were added to the cells every other day.

After five days in culture, the cells were trypsinized and the final cell density was determined using a Coulter® counter. (Coulter Electronics, Hialeah, FL, USA). The cells were released from the plates by replacing the culture medium with a solution containing 0.9% NaCl, 0.01M sodium phosphate (pH 7.4), 0.05% trypsin, and 0.02% EDTA (STV). The cells were incubated in this solution for 5-10 minutes at 37°C, and then the stock culture medium was added to the cells. The cells were then counted using a Coulter® counter.



Results shown in Figure 9 demonstrates the effect of different dilutions of FGF 98 on granulosa cells as compared to basic FGF. The figure legend for Figure 9 is as follows: O=basic FGF, =FGF 98.

The final cell density was graphed as a function of protein concentration. The protein concentration is graphed on a log scale. The protein concentration was determined by Bradford assay according to instructions accompanying the protein assay kit from BIORAD (Richmond, CA, USA).

The  $ED_{50}$  was calculated by (a), dividing in half the difference between the lowest and highest cell density value of the curve; and (b), determining from the graph what protein concentration corresponds to that cell density number obtained in (a). According to Figure 9 the  $ED_{50}$  for bFGF was found to be approximately 40 pg/ml, that for FGF 98 was about 100 ng/ml. The results of this assay demonstrate that FGF 98 was about 2500-fold less active than bFGF.

#### EXAMPLE 4

##### ABSENCE OF FGF 98 BIOACTIVITY ON ABAE OR ACE CELLS

FGF 98 can be characterized by its lack of activity on vascular endothelial cells derived from large vessels ("ABAE") or capillary cells ("ACE") as compared with basic FGF ("bFGF"). Stock cultures of ABAE and ACE cells were grown and maintained in Dulbecco's modified Eagle medium supplemented with 10% bovine serum, 0.25  $\mu$ l/ml fungizone, and 2 ng/ml bFGF. The cells were incubated at 37°C with a 10% CO<sub>2</sub> concentration and 99% humidity.

In the mitogenic assay, either 10<sup>4</sup> ABAE or 5x10<sup>3</sup> ACE cells were plated per well in 12-well plates in stock culture medium, as described in Gospodarowicz et al., *Proc. Natl. Acad. USA* 73:4120-4124 (1976), Gospodarowicz et al., *J. Cell. Physiol.* 127:121-136 (1976); and Gospodarowicz et al., *Proc. Natl. Acad. USA* 86:7311-7315 (1989). Ten microliter aliquots of the fractions to be tested were diluted into 1 ml of 0.2% (w/v) gelatin in phosphate buffered saline. Ten microliters of this dilution were added to cells seeded in 12-well cluster plates. A 10  $\mu$ l aliquot of either the diluted purified FGF 98 medium-containing fractions or 2 ng bFGF was added every other day.

After five days in culture, the cells were trypsinized and the final cell density was determined using a Coulter® counter. The cells were released from the plates by replacing the medium with a solution containing 0.9% NaCl, 0.01%M sodium phosphate (pH 7.4), 0.05% trypsin, and 0.02% EDTA (STV). The cells were incubated in this solution for 5-10 minutes at 37°C, and then the stock culture medium was added to the cells. The cells then were counted using a Coulter® counter (Coulter Electronics, Hialeah, FL, USA).

In the absence of heparin, results demonstrated the lack of activity of FGF 98 on ABAE and ACE cells in contrast to bFGF. In the presence of heparin (2 µg/ml), FGF 98 was bioactive on ABAE as well as ACE cells.

### EXAMPLE 5

#### FGF 98 EXPRESSION PATTERNS IN EMBRYOGENESIS

To determine the expression pattern of FGF 98 during embryogenesis, wholemount in situ hybridizations were performed on mouse embryos ranging in stage from embryonic day (E) 7.5 to E11.5 and on limbs from E12.5 to E14.5. For wholemount RNA in situ hybridizations, mouse embryos (CD-1) were isolated, fixed, and processed following the protocol by Nieto, M.A., Patel, K., Wilkinson, D.G. 1995. In situ hybridization analysis of chick embryos in whole mount and tissue section. In: *Methods in Avian Embryology* (ed. M. Bronner-Fraser), San Diego, Academic Press. Antisense riboprobe for FGF 98 was synthesized using the cDNA corresponding to EST accession number AA656693 as a template, linearized with XbaI and transcribed with T7 RNA polymerase. This probe encompasses C-terminal coding sequence and 3' untranslated region. The *fgf-8* probe was prepared as described by Crossley et al., *Development* 121:439-451 (1995). Embryos processed for wholemount in situ hybridization were equilibrated in 30% sucrose, embedded in OCT or 15% sucrose, 7.5% gelatin and cryosectioned at 12 or 30 µm, respectively.

At E7.5, no FGF 98 transcripts were detectable. At E8.5, FGF 98 transcripts were present in the paraxial mesoderm and in the newly formed somites. At all subsequent stages analyzed, FGF 98 continued to be expressed in the rostral part of the paraxial mesoderm and throughout the newly formed somites. As the more rostral somites mature,

FGF 98 transcripts first become restricted to the caudal part of the somite, and then to the caudal part of the myotome.

In the developing limbs, FGF 98 expression was initiated at E10.5 in a dorsal and a ventral domain in the mesenchyme. At E11, two separate domains of expression were observed on the ventral side of the limb. From E11.5 to at least E14.5, FGF 98 expression appeared to outline the differentiating skeletal elements, first the humerus, radius and ulna, followed by the digits. Transverse sections through E13.5 digits demonstrated that FGF 98 expression was present in tissues surrounding the differentiating cartilage elements. Furthermore, FGF was expressed in the interdigital regions at E12.5.

In the developing nervous system, FGF 98 transcripts were detected from E9.0 through at least E11.5. From E9.0 to E9.5, FGF 98 was transiently expressed in the caudal part of the neural plate which has not yet closed. In the differentiating neural tube, FGF 98 expression was initiated at E 9.5, in a dorsal and a ventral domain in the mantle layer. Furthermore, FGF 98 expression was observed in the dorsal root ganglia and in a subset of cranial ganglia (v, vii-viii, ix-x).

During the early development of the brain, the expression pattern of FGF 98 resembled that of *fgf-8*. At E8.5, FGF 98 expression was restricted to a narrow stripe in the head similar to a domain observed for *fgf-8* transcripts (Crossley et al., *Development* 121:439-451 (1995)). At subsequent stages, this FGF 98 expression domain marked the border between the midbrain and the hindbrain. However, the expression domain of FGF 98 in this region appeared to be wider than that of *fgf-8*. In the commissural plate of the telencephalon, FGF 98 expression was initiated at E9.0, approximately 0.5 days later than that of *fgf-8*. In the diencephalon, FGF 98 transcripts were first detected at E9.5 in both a dorsal and a ventral domain, and by E10.5, dorsal expression extended from the diencephalon rostrally into the telencephalon. In the optic stalks, FGF 98 expression was restricted ventrally, similar to *fgf-8*.

Finally, FGF 98 mRNA was detected in the pharyngeal region beginning at E9.5 in a restricted region of the second and the third branchial arch.

## EXAMPLE 6

## STEM CELL PROLIFERATION ASSAY

The neuron stem cell/progenitor assay involved the culturing of hippocampal cells from the embryonic rat brain for 6 days at 37°C in medium containing FGF 98, basic FGF, or no added factor. Neuronal survival and proliferation were assessed visually under a microscope on day 6. Photomicrographs were made and the cell number in each condition counted (the data are shown in Figure 10; series 1, 6 days in vitro DIV). The cultures were then washed and fresh media added without added FGF 98 or bFGF. Following 15 days in media without added factors cells were visually inspected, photomicrographs made and cells counted (Figure 11, series 2, 21 DIV). FGF 98 increases the number of neural cells at both 6DIV and 21DIV. FGF 98 stimulates the proliferation of neural progenitors under these conditions. The effects of FGF 98 are significantly different than the effect of bFGF where there is proliferation at 6 DIV but many of these cells die following withdraw of the growth factor. Few cells die in the FGF 98 treatment conditions.

Based on these and other results, FGF 98 can be used for the *ex vivo* expansion of precursor cells on the nervous system. Transplant or engraftment of cells can be used as a therapy for diseases in which specific populations of neurons are lost due to degeneration such as, for example, Parkinson's disease and Alzheimer's disease.

## EXAMPLE 7

## PREPARATION OF PRIMARY HIPPOCAMPAL CELL CULTURES

Hippocampi were dissected from E18-19 rats. The tissue was collected in Hibernate™ media (Gibco BRL) and minced into approximately 1mm pieces. Tissue was dissociated using the Papain Dissociation System (Worthington Biochemical Corporation). Following isolation the cells were resuspended in serum-free media composed of Neurobasal™ (Gibco BRL), 2%B27 supplement (GibcoBRL), L-glutamate and antibiotics. At the time of plating the growth factors were added: bFGF at 10ng/ml or FGF 98 at 200ng/ml. Cells were plated in 35mm tissue culture dishes coated with poly-L-lysine at a concentration of  $7.5 \times 10^4$  cells per dish. Following 6 days at 37°C in 5%CO<sub>2</sub> cells were rinsed

and fed with fresh media; no growth factors were added. Fifteen days after the factors were removed (total 21 DIV) the cells were fixed for 30 min in 4% paraformaldehyde and photographed. Some cultures are stained with the monoclonal antibody to the nestin intermediate filament (Developmental Studies Hybridoma Bank). This antibody recognizes neural stem cells/precursors. (Johe, K.K., et al. (1996). *Genes and Development* 10:3129-3140.) Figure 11 shows the expression of nestin in 15 DIV cultures treated with either bFGF or FGF 98 for the first 6 days. There are many more nestin positive cells in the FGF 98 treated cultures. See Figure 11, Panel A, nestin expression in bFGF treated cultures, and Figure 11, Panel B, nestin expression in FGF 98 treated cultures.

### EXAMPLE 8

#### FGF 98 EXPRESSION

A survey of FGF 98 expression was performed in mouse embryonic and adult tissues (>3months) using in situ hybridization. FGF 98 is expressed in the E14 mouse nervous system; areas of expression include the motor neurons of the brainstem and spinal cord, the dorsal root and cranial ganglia of the peripheral nervous system, the ventral tegmentum, developing forebrain, the region of the isthmus and developing cerebellum. In the adult brain FGF 98 is expressed in both the brain and spinal cord including the motor neurons, the substantia nigra, the deep cerebellar nuclei, the red nucleus and other areas. There are also several non-neuronal tissues in which FGF 98 is expressed including lung, heart, developing bone and chondrocytes.

Figure 12, Panel A, shows *in situ* hybridization of the adult brain stem and cerebellum. FGF 98 mRNA expression in deep cerebellar nuclei, and motor neurons of the facial and cochlear nuclei is visible. Figure 12, Panel B shows a high magnification of FGF 98 expression in motor neurons of the facial nucleus.

## EXAMPLE 9

PREPARATION OF ANTISERA TO FGF 98 BY IMMUNIZATION OF RABBITS  
WITH AN FGF 98 PEPTIDE.

The peptide sequence corresponding to amino acids 176-190 of the human FGF 98 protein was synthesized and coupled to keyhole limpet hemocyanin (KLH) as described (Harlow and Land, Antibodies: A Laboratory Manual, 1988. Cold Spring Harbor Laboratory, New York, NY) The KLH-coupled peptide was submitted to EL Labs and each of three rabbits were immunized. (EL Labs, Soquel, CA).

Peptide sequence: H-KRYPKGQPELQKPFFK-OH (KLH) peptide, synthesized by Research Genetics, Huntsville, AL.

Deposit Information:

The following materials were deposited with the American Type Culture Collection:

<u>Name</u>	<u>Deposit Date</u>	<u>Accession No.</u>
<i>Escherichia coli</i> 5 $\alpha$ FGF98	February 26, 1998	98662
<i>Escherichia coli</i> DH5- $\alpha$ FGF8h#4	February 18, 1998	98677

The above materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, VA 20110-2209, telephone 703-365-2700, under the accession numbers indicated. This deposit will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for purposes of Patent Procedure. The deposits will be maintained for a period of 30 years following issuance of this patent, or for the enforceable life of the patent, whichever is greater. Upon issuance of the patent, the deposits will be available to the public from the ATCC without restriction.

These deposits are provided merely as convenience to those of skill in the art, and are not an admission that a deposit is required under 35 U.S.C. § 112. The sequence of the polynucleotides contained within the deposited materials, as well as the amino acid

1000

2

2

[illegible]

## POLYNUCLEOTIDE AND POLYPEPTIDE SEQUENCES.

SEQ ID NO:1 is the nucleotide sequence.

The reading frame starts at nucleotide 610

CTATAGAAGGTACGCTGCAGGTACCGGTCCGGAATTCCTGGGTCGACCCACGCGTCCGCCACGCGTCCGCGGACGCGTGGGGACGCG  
GAGGAGGAGACATGAGCCGGCGGGCGCCAGACGGAGCGGCGTGACGCTTTCGCGCTGCAGCCGCGCGCCCGACCCCGAGCGCTGA  
CCCCTGCCCCACGACGCTCCGCGCCGGGCGGAGAGCGCAACTCGGCTTCCAGACCCGCGCGCATGCTGTCCCGGACTGAGCCGG  
GCAGCCAGCCTCCACGGACGCCCCGACGGCGGGCGGCCAGCAGTGAGCGAGCTTCCCGCACCGGCCAGGCGCTTCTGCACAACGG  
CTGCGCCCCCGACGCCCCGCGCCAGCCGGAGGGCGCAGCGCTCGGGAGGAGCGCGCGGGGCGCTGATGCCGACGGGCGCGCGCGG  
AGCGCCCCGGAGCAGCAGAGTCTGCAGCAGCAGCAGCGGCGAGGAGGAGCAGCAGCAGCGGCGGCGGCGGCGGCGGCGGCGGAG  
GCGCCCCGGTCCCGGCGCGCGGAGCGGACATGTGACGGCTGGGCTAGGAGCGCGCGCTCCCTCCCGCCAGCGATGTATTAGCGCCC  
TCCGCTGCACCTTGCTGTGTTTACACTTCTGTGCTGTGCTTCCAGGTACAGGTGCTGGTTGCCGAGGAGAAGCTGGACTTCCGAT  
CCAGTGGAGAACCAGACGCGGGCTCGGACGATGTGAGCCGTAAGCAGCTGCGGCTGTACCAGCTCTACAGCCGACAGTGGGAAAC  
ACATCCAGGTCTTGGGCGCAGGATCAGTGCCTCGCGGCGAGGATGGGACAAGTATGCCAGCTCCTAGTGGAGACAGACCTTCGGT  
AGTCAAGTCCGATCAAGGGCAAGGAGACGGAATTCACCTGTGCTGAACCGCAAAGGCAAGCTCGTGGGAAGCCGATGGCACCAG  
CAAGGAGTGTGTTCATCGAGAAGGTTCTGGAGAACAACACACGGCCCTGATGTCGGCTAAGTACTCCGGCTGGTACGTGGGCTTCA  
CCAAGAAGGGGCGGCGCTCTAGAGGATCCCCTGAGGGGCCCAAGCTTACGCGTGATGC

SEQ ID NO:2 is the amino acid sequence

MYAPSACTCLCLHFLLLCFQVQVLVAENVDFRIHVENQTRADDVSRQLRLYQLYSRTSGKHIQVLGRRISARGEDGDKYAQLLVE  
TDTFGSQVRIKGKETEFYLCMNRKGKLVGKPDGTSKECVFIEKVLNNY TALMSAKYSGWYVGFTKKGRPL

SEQ ID NO:3 is the primer sequence

CGGAGGATGGGGACAAGTAT

SEQ ID NO:4 is the FGF coding region with the native human 5' and 3' untranslated region

CCCACGCGTCCGCGGACGCGTGGGGACGCGGAGGAGAGACATGAGCCGGCGGGCGCCAGACGGAGCGG  
CCGTGACGCTTTCGCGCTGCAGCCGCGCGCCCCGACCCGGAGCGCTGACCCCTGGCCCCACGACAGCTCC  
GCGCCCGGGCGGAGAGCGCAACTCGGCTTCCAGACCCGCGCGCATGCTGTCCCGGACTGAGCCGGG  
AGCCAGCCTCCACGGACGCCCCGACGGCGGGCGGCCAGCAGTGAGCGAGCTTCCCGCACCGGCCAGG  
CGCTCTCTGCACAACGGCTGCGGCCCCGACGCCCTGCGCCAGCCGGAGGGCGCAGCGCTCGGGAGGAG  
CCGCGCGGGGCGCTGATGCCGAGGGCGCGCGGAGCGCCCCGGAGCAGCAGAGTCTGCAGCAGCAGC  
AGCCGGCGAGGAGGGAGCAGCAGCAGCGCGCGGCGGGCGGCGGCGGCGGCGGAGGCGCCCGGTCCCGG  
CGCGCGGAGCGGACATGTGACGGCTGGGCTAGGAGCGCGCGCTCCCTCCCGCCAGCGATGTATTAGC  
GCCCTCCGCTGCACCTTGCTGTGTTTACACTTCTGTGCTGTGCTTCCAGGTACAGGTGCTGGTTGCC  
GAGGAGAACGTGGACTTCCGATCCACGTGGAGAACCAGACGCGGGCTCGGACGATGTGAGCCGTAAGC



AGCTGCGGCTGTACCAGCTCTACAGCCGGACCAGTGGGAAACATCCAGGTCCTGGGCCGAGGATCAG  
 TGCCCGCGGCAGGATGGGGACAAGTATGCCAGCTCCTAGTGGAGACAGACACCTTCGGTAGTCAAGTC  
 CGGATCAAGGGCAAGGAGACGGAAATCTACCTGTGCATGAACCGCAAAGGCAAGCTCGTGGGGAAGCCCG  
 ATGGCACCAGCAAGGAGTGTGTTCATCGAGAAGGTTCTGGAGAACAACACACGGCCCTGATGTCGGC  
 TAAGTACTCCGGCTGGTACGTGGGCTTCACCAAGAAGGGCGGCCGGAAGGGCCCCAAGACCCGGGAG  
 AACCAGCAGGACGTGCATTTTCATGAAGCGCTACCCCAAGGGGCAGCCGGAGCTTCAGAAGCCCTTCAAGT  
 ACACGACGGTGACCAAGAGGTCCCGTCGGATCCGGCCACACACCTGCCTAGGCCACCCCGCCGCGGCC  
 CCTCAGGTCGCCC TGCCACACTCACACTCCCAGAAAACGTCATCAGAGGAATATTTTACATGAAAAAT  
 AAGGAAGAAGCTCTATTTTTGTACATTGTGTTTAAAGAAGACAAAACTGAACAAAACCTTGGGGGG  
 AGGGGTGATAAGGATTTTATTGTTGACTTGAAACCCCGATGACAAAAGACTCACGCAAAGGGACTGTAG  
 TCAACCCACAGGTGCTTGTCTCTCTAGGAACAGACAACCTAAACTCGTCCCAGAGGAGGACTTGAA  
 TGAGGAAACCAACACTTTGAGAAGCCAAAGTCTTTTTCCAAAGGTTCTGAAAGGAAAAAAAAAAC  
 AAAAAAAAAAAAAAAAAAAAAAAAAA

SEQ ID NO: 5 is the full length amino acid sequence of FGF

MYSAPSACTCLCLHFLLLCFQVQVLVAENVDFRIHVENQTRARDDVSRKQLRLYQLYSRTSGKHIQVLG  
 RRISARGEDGDKYAQLLVETDTFGSQVRIKGKETEFYLCMNRKGKLVGKPDGTSKECVFIEKVLNNYTA  
 LMSAKYSGWYVGFTKKGRPRKGPKTRENQDVFHMKRYPKQPELQKPFKYTTVTKRSRRIRPTHPA

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118